

Influencing a Particle

L'Architech (Gabriel Breton-Breault)

Introduction

Influencing a particle is never an isolated phenomenon: it is a directional event that transforms its internal structure. In directional cosmology, each particle possesses its own vibration, spin, directional pressure, and geometric orientation that define its state. Thus, any interaction—repulsion, confinement, energy variation, or photon trapping—modifies the very nature of the particle.

When two particles repel each other, this repulsion generates tension within their directional fields. This tension produces an internal variation, a measurable energetic event that alters the vibration of the confined strings. Similarly, trapping a photon amounts to imposing a geometry upon it: the free string becomes a confined string, and the particle changes its nature.

In an entangled system, influencing a particle through an attosecond-synchronized laser pulse creates a directional event. The linked particle may reflect this variation in its own state, not as a classical transmission of information, but as a directional resonance between two correlated structures.

This document explores these phenomena precisely: how a particle can be influenced, how its nature can change, and how these directional events become readable within an architecture such as the VSMD.

Chapter 1 — Directional Transformation of Particles

Influencing a particle means modifying the internal geometry of its vibration. In directional theory, no particle is static: each is defined by a confined or free string, a directional pressure, a spin, and a geometric orientation that evolve according to interactions. Thus, any external modification—repulsion, confinement, energy variation, or photon trapping—transforms the nature of the particle.

1.1 — Repulsion as a Directional Event

When two particles repel each other, the observed force is only the outward manifestation of a deeper phenomenon. Repulsion creates directional tension, a compression of their internal fields. This compression modifies the vibration of the confined strings, producing a directional event: a measurable internal variation that alters spin, pressure, and orientation.

In this framework, repulsion is not a simple mechanical interaction. It is an energetic signal, an internal transformation that can be interpreted within a directional system such as the VSMD.

1.2 — Trapping a Photon: Changing the Nature of the Particle

A free photon is a pure directional string, without spin, confinement, or imposed orientation. But when a photon is trapped, its geometry changes: the free string becomes a confined string, and the particle ceases to be a photon in the directional sense.

Confinement imposes an orientation, modifies the vibration, and creates a new directional state. Thus, trapping a photon amounts to changing its nature, because the particle acquires properties it did not possess in its free state.

This transformation is essential for understanding how an energetic event can be generated, read, and interpreted within a directional architecture.

1.3 — Entanglement and Directional Influence

In an entangled system, two particles share a correlated vibrational structure. Influencing one of them—through repulsion, energy variation, or laser pulse—creates a directional event in its confined string. The linked particle may reflect this variation in its own state, not as a classical transmission of information, but as a directional resonance.

For this resonance to be readable, the laser pulses must be synchronized at the attosecond scale, ensuring an absolute temporal reference. This synchronization makes it possible to distinguish a real variation from measurement noise, and to transform the directional event into an interpretable signal.

1.4 — Bringing Two Entangled Systems Closer

When two entangled particles are brought near another entangled pair, their directional fields enter into resonance. This resonance can influence the internal vibration of all four particles, creating a series of directional events. In this framework, proximity is not a simple spatial closeness: it is a geometric interaction between two coherent vibrational structures.

Chapter 2 — The Directional Event

A directional event is the internal transformation of a particle caused by an external variation. Unlike classical physics, where interaction is described as a force or energy transfer, directional theory considers each interaction as a modification of the particle's vibrational geometry. This modification is not a secondary effect: it constitutes the event itself.

2.1 — Definition of a Directional Event

A directional event occurs when a particle undergoes a variation in:

- its internal vibration,
- its directional pressure,
- its spin,
- or its geometric orientation.

These variations are not independent: they are linked to the structure of the confined or free string that composes the particle. Thus, any change in the directional environment—repulsion, confinement, photon trapping, laser pulse—modifies the string, and therefore the particle.

A directional event is therefore a measurable internal transformation, which can be interpreted within an architecture such as the VSMD.

2.2 — Origin of Events: Repulsion and Directional Tension

When two particles repel each other, the observed force is only the surface of the phenomenon. Repulsion creates tension within their directional fields, which modifies the internal vibration of the confined strings. This modification is a directional event.

In this framework, repulsion is not a simple displacement: it is an internal reorganization, a vibrational response that changes the nature of the particle.

2.3 — Photon Trapping: From Free String to Confined String

Trapping a photon amounts to imposing a geometry upon it. The free photon, which has neither spin nor imposed orientation, becomes a confined string. This transition changes its vibration, its directional pressure, and its behavior.

Thus, photon trapping is a major directional event: it transforms a luminous particle into a geometrically constrained particle.

This transformation is essential for understanding how a photon can become an active element in a directional architecture.

2.4 — Influence Through Laser Pulse

Increasing the intensity of a laser on an entangled particle modifies its directional pressure. This modification creates an internal event, which can be detected if the laser pulses are synchronized at the attosecond scale.

Extreme synchronization makes it possible to distinguish:

- a real variation,
- from measurement noise,
- or thermal fluctuation.

In an entangled system, this event may be reflected in the linked particle, not as a classical transmission of information, but as a directional resonance.

2.5 — Interaction Between Entangled Systems

When two entangled particles are brought near another entangled pair, their directional fields enter into resonance. This resonance can influence the internal vibration of all four particles, creating a series of directional events.

In this framework, proximity is not a simple spatial closeness: it is a geometric interaction between two coherent vibrational structures.

Chapter 3 — Directional Reading

Directional reading is the process by which an internal event generated within a particle becomes detectable and interpretable. In directional theory, reading a particle does not consist of measuring a fixed value, but of observing a **geometric variation** in its vibration, spin, or directional pressure. This variation is the imprint of the directional event.

3.1 — Reading a Particle: A Geometry, Not a Value

In classical physics, a measurement is a value: frequency, energy, spin, polarization. In directional cosmology, a measurement is a vibrational geometry.

Reading a particle means analyzing:

- the shape of its vibration,
- the tension of its confined string,
- its internal directional pressure,
- and the geometric orientation imposed by the environment.

Thus, directional reading is not a snapshot: it is a dynamic interpretation of internal transformation.

3.2 — The Importance of Attosecond Synchronization

For reading to be reliable, laser pulses must be synchronized at the attosecond scale. This extreme synchronization allows:

- establishing an absolute temporal reference,
- distinguishing real variation from measurement noise,
- isolating thermal fluctuations,
- and detecting directional events with precision.

Without this synchronization, directional reading would be drowned in quantum noise. With it, each variation becomes an interpretable signal.

3.3 — Reading a Directional Event

When a particle undergoes a directional event—repulsion, photon trapping, energy variation—its internal vibration changes. This variation can be detected through:

- a modification of spin,
- tension in the confined string,
- variation in directional pressure,
- or alteration of geometric orientation.

Directional reading consists of identifying this variation, then interpreting it within the VSMD architecture.

3.4 — Reading in an Entangled System

In an entangled system, two particles share a correlated vibrational structure. Influencing one of them creates a directional event in its confined string. The linked particle may reflect this variation in its own state.

Directional reading then makes it possible to detect:

- the variation in the first particle,
- the resonance in the second,
- and the coherence between the two events.

This is not a classical transmission of information. It is a reading of directional resonance.

3.5 — Reading Between Nearby Entangled Systems

When two entangled particles are brought near another entangled pair, their directional fields enter into resonance. This resonance can generate a series of directional events in all four particles.

Directional reading makes it possible to observe:

- geometric influence between systems,
- propagation of variation,
- and global vibrational coherence.

In this framework, reading is not a simple measurement: it is a geometric analysis of a complex directional system.

Conclusion

Directional theory establishes a new way of understanding particle interaction. In this framework, a particle is never an isolated point: it is a vibrational structure, a confined or free string defined by its directional pressure, spin, and geometric orientation. Any interaction—repulsion, photon trapping, energy variation, or laser influence—becomes a directional event, an internal transformation that changes the nature of the particle.

Entanglement, far from being a classical communication channel, constitutes vibrational coherence between two particles. Influencing one of them creates a directional event that may be reflected in the linked particle, not as a transmission of information, but as geometric resonance. This resonance becomes readable thanks to attosecond-synchronized laser monitoring, which provides an absolute clock capable of isolating each internal variation.

The VSMD architecture relies entirely on this logic: communication is not wave propagation, but interpretation of directional events. A signal is not sent: it is detected, read, and interpreted within the vibrational geometry of entangled particles. Thus, the VSMD communicates without frequency, without waves, without classical transmission—only through the reading of internal variations within a coherent directional structure.

This document establishes the foundations of this architecture: particle transformation, the nature of directional events, geometric reading, and communication through resonance. It opens the way to a

physics in which the internal geometry of particles becomes the language, and directionality replaces propagation.

Directional communication in the VSMD is not a technology: it is a **natural consequence of the vibrational structure of the universe**.

UPDATE — Reference Particles and Directional Timeline Analysis

This update introduces a fundamental concept for directional communication: **the use of entangled reference particles** and **the analysis of the reading timeline** as the foundation of an interpretable directional language.

1 — Entangled Reference Particles

In classical approaches, an entangled particle is measured in isolation, making any attempt at communication impossible: its individual state is random. Directional theory resolves this limitation by introducing a **pair of entangled reference particles**, maintained in a stable state and monitored by attosecond-synchronized laser pulses.

This reference pair serves as a **baseline directional geometry**, allowing the system to distinguish:

- natural variations,
- thermal fluctuations,
- genuine directional events.

It constitutes the “directional zero,” the essential comparison state for any coherent reading.

2 — The Influenced Pair: Generating Events

A second entangled pair is used as the **directional channel**. By influencing one of the particles (repulsion, energy variation, photon trapping, laser pulse), a **directional event** is generated within its confined string.

This event is not a transmitted message: it is a **geometric variation** that becomes readable only when compared to the reference pair.

3 — Directional Timeline Analysis

Directional reading is not performed on a single value, but on a **sequence of measurements over time**. Each laser pulse becomes a point on a directional timeline.

By analyzing this timeline, one can detect:

- ruptures,
- recurring motifs,
- vibrational sequences,

- directional patterns.

These patterns are not bits: they are **temporal structures of variation**, impossible to obtain from an isolated reading.

The timeline thus becomes a **language space**, where directional events form interpretable signatures.

4 — Comparing Timelines: Birth of the Directional Signal

The signal never appears in the influenced particle alone. It emerges from the **difference** between:

- the timeline of the reference pair,
- the timeline of the influenced pair.

This comparison reveals:

- the intensity of the event,
- its duration,
- its internal geometry,
- its vibrational coherence.

This difference constitutes the directional information.

5 — Consequence: Communication Without Transmission

Thanks to reference particles and timeline analysis:

- no wave is sent,
- no frequency is propagated,
- no classical transmission occurs.

Directional communication relies solely on:

the reading of internal variations within a coherent entangled structure.

This mechanism fully respects standard physics while opening the way to a directional language based on temporal patterns.